

# Quantification of fiber production by preadipocytes



Adipose Tissues • 2D Cell Models • High Content Screening • Automatic Detection • Dermatology



## YOUR NEEDS

- Monitor pre-adipocytes differentiation
- Monitor the fibro-inflammation process (collagen, fibronectin)

## OUR SOLUTIONS

- Fully automatic imaging and image analysis
- Robust data, short delay and cost-saving

## General Procedure




### Image acquisition:

- Acquisition done with structured light or confocal microscopy
- Several fields of view to maximize the amount of data
- Image stack for each field of view to get each cellular structure on their focal plane


### Image processing:

- ① Detail of an image after acquisition with spinning disk confocal microscopy
- ② Denoising and segmentation of every fiber
- ③ Refining the segmented object and calculating the criteria: **total fiber length, thickness of fibers**
- ④ Automatic generation of illustrative images for every field of view



## Application example

During weight gain, adipose tissue is characterized by the presence of low-grade inflammation and matrix remodelling. This leads to a greater tissue stiffness and to resistance to weight loss.



©Imactiv-3D, Centre Pierre Potier, 1 place Pierre Potier, 31106 Toulouse, France • +33 5 82 99 10 09

[contact@imactiv-3d.com](mailto:contact@imactiv-3d.com) • [www.imactiv3d.com](http://www.imactiv3d.com)